FIELD DAY REPORT - 1992

Texas A&M University Agricultural Research and Extension Center at Overton

Texas Agricultural Experiment Station Texas Agricultural Extension Service

Overton, Texas

April 30, 1992

Research Center Technical Report 92-1

All Programs and information of the Texas Agricultural Experiment Station and Texas Agricultural Extension Service are available to everyone without regard to race, color, religion, sex, age, or national origin.

Mention of trademark or a proprietary product does not constitute a guarantee or a warranty of the product by the Texas Agricultural Experiment Station or Texas Agricultural Extension Service and does not imply its approval to the exclusion of other products that also may be suitable.

RABBITEYE BLUEBERRY PLANT RESPONSE TO NITROGEN AND PHOSPHORUS

V. A. Haby, J. V. Davis, and A. T. Leonard

Background. Acid, sandy soils selected for blueberry production are usually deficient in plant available phosphorus (P). Fertilizer addition at planting has not been a recommended practice. This glasshouse research was designed to study response of transplanted rabbiteye blueberry plants to fertilizer P and N rates.

Research Findings. Preplant phosphate was incorporated into 9.7 lb of 1:1 v/v loamy sand:peat mixture in a 2-gallon container at rates of 0, 41, 82, 123 and 164 lb P₂O₅/ac. Nitrogen was split-applied post-plant four times at 60-day intervals beginning 30 days after blueberry bushes were transplanted. Total N rates were equivalent to 0, 100, 200, or 300 lb/ac. Batches of 194 lb of the sand-peat mixture were treated with 37.8 gm of a fertilizer blend that contained 16% K₂O, 9.4% Ca, 8% Mg, 8% S, 1% Zn, 1% Fe, 0.2% Cu, 0.03% B, and 0.01% Mo. Plants were harvested and separated into leaf, stem, and root components 270 days after transplanting.

Total plant growth increased as N rate increased (Table 1). Leaf and stem weight increases contributed to this linear response while root weight was not affected by fertilizer N.

Table 1. Response of 'Tifblue' rabbiteye blueberry plants to fertilizer nitrogen and phosphorus.

				Total	Number
Nitrogen	Leaf	Stem	Root	Plant	Stems
rate	D.M. ^y	D.M.	D.M.	D.M.	>2 mm
lb/ac	grams/pot				
0	10.2	20.5	26.5	57.2	3.0
100	14.0	23.4	30.1	67.2	3.8
200	17.0	26.2	25.1	69.7	5.1
300	21.4	30.7	30.3	82.4	4.8
Linear	*** ^Z	***	N.S.	***	***
Quadratic	N.S.	N.S.	N.S.	N.S.	N.S.
P ₂ O ₅ rate					
lb/ac			,		
0	13.2	23.3	27.2	63.7	3.3
41	11.2	20.3	23.9	55.4	3.2
82	10.3	19.6	19.1	48.5	3.4
123	23.0	32.3	38.5	95.7	6.3
164	20.6	30.5	31.2	82.4	4.7
Linear	***	***	**	***	***
Quadratic	***	**	N.S.	*	N.S.

^yDry Matter

²***, **, and * indicate statistically significant linear or quadratic responses at p=0.01, 0.05, and 0.10, respectively, due to treatment.

Phosphorus at the 123 and 164 lb/ac rates increased total plant growth. Phosphorus applications increased leaf P levels, but did not increase soil test P (data not shown).

The greatest yield response to the N and P interaction occurred with plant top growth (Fig. 1). Except




FIGURE 1. Rabbiteye Blueberry Plant Response to the interactions of Nitrogen and Phosphorus

for the high N rate at the zero P treatment, N had no effect on plant growth at the 0, 41, and 82 lb P_2O_5 /acre application rates. There was a significant (p = 0.01) linear response to N at P_2O_5 rates of 123 and 164 lb/ac. The optimum P_2O_5 rate for fertilization of the soil:peat mixture in the planting hole under conditions of this experiment was the equivalent of 123 lb/acre.

Application. Results from this research indicated that a rate of P_2O_5 equivalent to 123 lb/ac was needed to increase rabbiteye blueberry plant growth when the soil pH is between 5.0 and 5.5. This rate equates to 5.1 gm 0-46-0 per 50 lb soil:peat mix (45 lb soil:5 lb peat). This amounts to approximately 0.2 oz (1 slightly rounded teaspoon) 0-46-0 added to 45 lb soil:5 lb peat, equivalent to three-fourths bushel to be placed into the planting hole. At pH below 5.0, increased levels of extractable Al occur in this potting medium and a higher rate of P may be needed for best plant growth.