

FRUIT AND NUT CROPS RESEARCH IN TEXAS, 1987

Page	Participating Scientists	Crops	Location of Research Station, and Cooperative Research Sites
3, 5	David H. Byrne	Peach	College Station
3, 5	Terry Bacon	Plums	
7	J. Dan Hanna	Apricots	
9	Calvin G. Lyons	Grapes	
10	T. Glynn Littleton	Pecans	
10	G. R. McEachern		
10, 20, 46	Gerald Johnson		
12	J. Benton Story		
46	Berry Thompson		
15	R. D. Mardard	Pecan	
17	L. Austin Stockton	Grapes	Fort Stockton
19, 20, 21, 23	John A. Lipe	Peach	Fredericksburg
19, 20	Duery Menzies	Pecan	

COMPILED AND EDITED BY:

Robert E. Rouse
Texas Agricultural Experiment Station
2415 East Highway 83
Weslaco, TX 78596

David H. Byrne
Department of Horticulture
Texas A&M University
College Station, TX 77843

The Texas Agricultural Experiment Station, Neville P. Clarke, Director,
Texas A&M University System, College Station, TX.

SUBJECT TOPIC: Chemical Mowing of Orchard Middles With Roundup, Fusilade and Poast

INVESTIGATOR(S): John A. Lipe - TAEX, Fredericksburg

CROP(S): Pecans

ABSTRACT:

Objectives:

1. Compare varying rates of Roundup herbicide and Roundup plus additives to determine optimum effects.
2. Compare chemical mowing effects of Roundup, Fusilade and Poast.

General Approach:

Initial chemical treatments were made on June 6, 1986, one week after mowing the predominantly Coastal bermudagrass orchard floor. Three of the treatments with Roundup were repeated on July 30.

Treatments were applied at 6.4 Km/hr (4 mph), 140 KPa (20 PSI), 84 l of solution per hectare (9 gal/acre) with a tractor-drawn sprayer with 80015 flat fan spray tips. Single plots 30' wide and 150' long were used for each treatment.

Findings:

Roundup at 0.9 l/ha (12 oz/acre) gave very satisfactory suppression of Bermudagrass growth (Table 1). Addition of 1% Uran (32-0-0) improved the suppression by Roundup.

Half rates of both Fusilade and Poast gave very good suppression of Bermudagrass.

Retreatment of the Roundup plots at 0.9 l/ha (12 oz. rates only) on July 30 gave less dramatic suppression, but were adequate to avoid the need for mechanical mowing until harvest.

A cost comparison of mechanical mowing at 2 mowings each of \$12.30/ha (\$5/acre) with each chemical mowing at \$4.92/ha (\$2/acre plus cost of chemical) produced the following comparison.

Roundup 0.36 l (12 oz)	-----	\$ 9.50
Fusilade 4E 0.12 l (4 oz)	---	9.80
Poast 0.36 l (12 oz)	-----	11.85
Mechanical Mowing	-----	10.00

Table 1. Chemical control of bermudagrass using selected herbicides.

Treatment (6/13/86)	6/26/86			7/30/86		
	Vigor (1-10)	Grass Ht cm (inches)		Vigor (1-10)	Grass Ht cm (inches)	
Check	10	25.4	10.0	5	30.5	12
Roundup 0.9 l/ha (12 oz/acre)	4	6.4	2.5	4	15.2	6
Roundup 0.9 l/ha (12 oz/acre)	4	6.4	2.5	4	15.2	6
+ Bivert 0.45 l/ha (6 oz/acre)						
Roundup 0.9 l/ha (12 oz/acre)	3	5.1	2.0	3	10.2	4
+ Uran (1%)						
Roundup 0.45 l/ha (6 oz/acre)	6	10.2	4.0	7	30.5	12
Roundup 0.45 l/ha (6 oz/acre)	6	10.2	4.0	7	30.5	12
+ Bivert 0.23 l/ha (3 oz/acre)						
Roundup 0.45 l/ha (6 oz/acre)	6	10.2	4.0	7	30.5	12
+ Uran (1%)						
Fusilade 0.29 l/ha (1/4 pt)	2	6.4	2.5	3	10.2	4
+ Natural Oil 2.3 l/ha (1 qt/acre)						
Poast 0.86 l/ha (3/4 pt/ac)	2	6.4	2.5	3	12.7	5
+ Natural Oil 2.3 l/ha (1 qt/acre)						